

# EBP50 Rabbit mAb

货号: **AYM28972**

## 产品信息

反应	Human,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:200
理论分子量	22kDa/38kDa
实测分子量	50kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HepG2,293T,HT-29,MCF7,Mouse liver,Mouse kidney,Rat liver
细胞定位	Apical cell membrane,Cell projection,Cytoplasm,Endomembrane system,Peripheral membrane protein,filopodium,microvillus,ruffle
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein corresponding to Human EBP50.
------	--

## 靶点信息

研究背景	This gene encodes a sodium/hydrogen exchanger regulatory cofactor. The protein interacts with and regulates various proteins including the cystic fibrosis transmembrane conductance regulator and G-protein coupled receptors such as the beta2-adrenergic receptor and the parathyroid hormone 1 receptor. The protein also interacts with proteins that function as linkers between integral membrane and cytoskeletal proteins. The protein localizes to actin-rich structures including membrane ruffles, microvilli, and filopodia. Mutations in this gene result in hypophosphatemic nephrolithiasis/osteoporosis type 2, and loss of heterozygosity of this gene is implicated in breast cancer.
基因ID	9368

基因名	SLC9A3R1
Swiss	O14745
别名	EBP50

## 产品验证



Western blot analysis of EBP50 expressed in Rat liver,HepG2,293T using EBP50 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)