— ABLYBIO, Help Your Research

Histone H2B (mono methyl K116) Rabbit mAb

货号: AYM28974

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	**WB:** 1:500 - 1:2000 **IHC:** 1:50 - 1:200 **IF/ICC:** 1:50 - 1:200
理论分子量	13kDa
实测分子量	14kDa
形式	Liquid
保存条件	Store at -20℃. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,C2C12,C6
细胞定位	Chromosome,Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human Histone H2B (mono methyl K116).

靶点信息

研究背景	Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octa mer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The l inker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chr omatin into higher order structures. This gene encodes a replication-dependent histone that is a member of the histone H2B family, and generates two transcripts through the use of the conserved stem-loop ter mination motif, and the polyA addition motif. The protein has antibacterial and antifungal antimicrobial ac tivity.
基因ID	8349
基因名	HIST2H2BE
Swiss	Q16778 (https://www.uniprot.org/uniprotkb/Q16778/entry)
别名	Histone H2B (mono methyl K116),Histone H2B (mono methyl K116) Rabbit mAb,HIST2H2BE,H2B-clustere d histone 21,Histone H2B-GL105,Histone H2B.q,H2BFQ

产品验证



Western blot analysis of Histone H2B (mono methyl K116) expressed in HeLa,C2C12,C6 using Histone H2 B (mono methyl K116) Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5 000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanc ed Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (https://www.ablybio.cn//www.ablybio.cn)