

IRAK2 Rabbit mAb

货号: **AYM28978**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IP FC
推荐浓度	WB: 1:500 - 1:2000 IP: 1:20 - 1:50 FC: 1:20 - 1:50
理论分子量	69kDa
实测分子量	69kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-549,HeLa,HepG2,NCI-H460,Mouse liver,Mouse kidney,Rat lung
细胞定位	cytoplasm,cytosol,nucleus,plasma membrane
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human IRAK2.
序列	MACYIYQLPSWVLDLDCRNMDALSEWDWMEFASYVITDLTQLRKIKSMERVQGV SITRELLWWWGMRQATVQQLVDLL CRLELYRAAQIILNWKPAPEIRCPAFAFPDSVKPEKPLAASVRKAEDEQEEGQPVRMATFPGPGSSPARAHQPAFLQPPEED APHSLRSDLPTSSDSKDFSTSIPKQEKLLSLAGDSLFWSEADV VQATDDFNQNRKISQGT FADVYRGHRHGKPFVFKLLR ETACSSPGSIERFFQAE LQICLRCHPNVL

靶点信息

研究背景	IRAK2 encodes the interleukin-1 receptor-associated kinase 2, one of two putative serine/threonine kinases that become associated with the interleukin-1 receptor (IL1R) upon stimulation. IRAK2 is reported to participate in the IL1-induced upregulation of NF-kappaB.
------	---

基因ID	3656
基因名	IRAK2
Swiss	O43187
别名	IRAK2;IRAK-2

产品验证



Western blot analysis of IRAK2 expressed in A-549, HeLa, HepG2 using IRAK2 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn