

Frizzled 9 / CD349 Rabbit mAb

货号: **AYM28993**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	64kDa
实测分子量	64kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	TE-1,293T,HepG2,THP-1,U-251MG,Mouse heart,Mouse brain,Rat skeletal muscle
细胞定位	Cell membrane,Multi-pass membrane protein
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human Frizzled 9 / CD349.
序列	ERLNMDFWRLRATEQPAAAAAGPGGRRDCSLPGGSVPTVAVFMLKIFMSLVVGITSGVWVWSSKTFQTWQSLCYRKIAA GRARAKACRAPGSYGRGTHCHYKAPTIVLHMTKTDPSLENPTHL

靶点信息

研究背景	Members of the 'frizzled' gene family encode 7-transmembrane domain proteins that are receptors for Wnt signaling proteins. The FZD9 gene is located within the Williams syndrome common deletion region of chromosome 7, and heterozygous deletion of the FZD9 gene may contribute to the Williams syndrome phenotype. FZD9 is expressed predominantly in brain, testis, eye, skeletal muscle, and kidney.
基因ID	8326

基因名	FZD9
Swiss	O00144
别名	FZD9;CD349;FZD3

产品验证



Western blot analysis of Frizzled 9 / CD349 expressed in Mouse heart,Rat skeletal muscle,TE-1 using Frizzled 9 / CD349 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1: 5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn