

Fetuin A Rabbit mAb

货号: **AYM28996**

产品信息

| | |
|-------|---|
| 反应 | Human |
| 宿主 | Rabbit |
| 克隆性 | Monoclonal |
| 预测反应 | |
| 应用 | WB IP |
| 推荐浓度 | WB: 1:500 - 1:2000 IP: 1:20 - 1:50 |
| 理论分子量 | 39kDa |
| 实测分子量 | 58kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA, 50% glycerol, pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | 293T, HepG2, DU145, A-549 |
| 细胞定位 | Secreted |
| 纯化 | Affinity purification |

抗原信息


| | |
|------|--|
| 抗原信息 | Recombinant fusion protein corresponding to Human Fetuin A. |
| 序列 | APHGPGLIYRQPNCDDPETEEAALVAIDYINQNLPGWYKHTLNQIDEVKVWPQQPSGELFEIEIDTLETTCHVLDPTPVARC SVRQLKEHAVEGDCDFQLKLDGKFSVYAKCDSSPDSAEDVRKVCQDCPLLAPLNDTRVVHAAKAALAFNAQNNGS NFQLEEISRAQLVPLPPSTYVEFTVSGTDCVAKEATEAAKCNLLAEKQYGFCKATLSEKLGGAEVAVTCMVFTQPVSSQP QPEGANEAVPTPVDPDAPSPPLGAPGLPPAGSPDHSV |

靶点信息

| | |
|------|--|
| 研究背景 | Alpha2-HS glycoprotein (AHSG), a glycoprotein present in the serum, is synthesized by hepatocytes. The AHSG molecule consists of two polypeptide chains, which are both cleaved from a proprotein encoded from a single mRNA. It is involved in several functions, such as endocytosis, brain development and the formation of bone tissue. The protein is commonly present in the cortical plate of the immature cerebral cortex and bone marrow hemopoietic matrix, and it has therefore been postulated that it participates in the development of the tissues. However, its exact significance is still obscure. |
|------|--|

| | |
|-------|---|
| 基因ID | 197 |
| 基因名 | AHSG |
| Swiss | P02765 |
| 别名 | AHSG; A2HS; AHS; FETUA; HSGA; alpha-2-HS-glycoprotein |

产品验证



Western blot analysis of Fetuin A expressed in 293T, HepG2, DU145 using Fetuin A Rabbit mAb at 1:10 00. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn