

Jarid2 Rabbit mAb

货号: **AYM29008**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	106kDa/119kDa/138kDa
实测分子量	139kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	HL-60, Mouse thymus, Rat heart
细胞定位	Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human Jarid2.
序列	ETVHFATTQWTSMGFETAKEMKRRIHAKPFSMEKLLYQIAQAEAKKENGPTLSTISALLDELDRDTELRRQLFEAGLHSSA RYGSHDGSSTVADGKKKPRKWLQLETSEKRCQICQHLCYLSMVVQENENVVFCLECALRHVEKQKSCRGLKLMYRYDEE QIISLVNQICGKVSNGKNGSIENCLSKPTPKRGPRKRATVDVPPSRLSASSSSKSASSSS

靶点信息

研究背景	This gene encodes a Jumonji- and AT-rich interaction domain (ARID)-domain-containing protein. The encoded protein is a DNA-binding protein that functions as a transcriptional repressor. This protein interacts with the Polycomb repressive complex 2 (PRC2) which plays an essential role in regulating gene expression during embryonic development. This protein facilitates the recruitment of the PRC2 complex to target genes. Alternate splicing results in multiple transcript variants. Mutations in this gene are associated with chronic myeloid malignancies.
------	---

基因ID	3720
基因名	JARID2
Swiss	Q92833
别名	JARID2;JMJ

产品验证



Western blot analysis of Jarid2 expressed in HL-60, Mouse thymus, Rat heart using Jarid2 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn