

PHD2 / prolyl hydroxylase Rabbit mAb

货号: **AYM29010**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IP
推荐浓度	WB: 1:500 - 1:2000 IP: 1:20 - 1:50
理论分子量	36kDa/43kDa/46kDa
实测分子量	46kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	293T, A-549, Mouse heart
细胞定位	Cytoplasm, Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human PHD2 / prolyl hydroxylase.
序列	MANDSGGGPGGPPSPSERDRQYCELCGKMENTLLRCSRCSFSCYCKEHRQDQWKKHKLVCQSGSEGALGHGVGPHQHSG PAPPAAVPPPFRAGAREPRKAAARR

靶点信息

研究背景	The protein encoded by this gene catalyzes the post-translational formation of 4-hydroxyproline in hypoxia-inducible factor (HIF) alpha proteins. HIF is a transcriptional complex that plays a central role in mammalian oxygen homeostasis. This protein functions as a cellular oxygen sensor, and under normal oxygen concentration, modification by prolyl hydroxylation is a key regulatory event that targets HIF subunits for proteasomal destruction via the von Hippel-Lindau ubiquitylation complex. Mutations in this gene are associated with erythrocytosis familial type 3 (ECYT3).
------	--

基因ID	54583
基因名	EGLN1
Swiss	Q9GZT9
别名	EGLN1;C1orf12;ECYT3;HALAH;HIF-PH2;HIFPH2;HPH-2;HPH2;PHD2;SM20;ZMYND6

产品验证



Western blot analysis of PHD2 / prolyl hydroxylase expressed in 293T, A-549, Mouse heart using PHD2 / prolyl hydroxylase Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn