

KIF5B Rabbit mAb

货号: **AYM29016**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC IP
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 IP: 1:20 - 1:50
理论分子量	109kDa
实测分子量	120kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,Jurkat,HepG2,Mouse brain,Mouse liver,Mouse kidney,Rat brain
细胞定位	Cytoplasm,cytoskeleton
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human KIF5B.
序列	LQNVEQKKRQLEESVDALSEELVQLRAQEKVHEMEKEHLNKKVQTANEVKQAVEQQIQSHRETHQKQISSLRDEVEAKAKLI TDLQDQNQKMMLEQERLRVEHEKCLKATDQEKSRKRLHELTV

靶点信息

研究背景	Microtubule-dependent motor required for normal distribution of mitochondria and lysosomes. Can induce formation of neurite-like membrane protrusions in non-neuronal cells in a ZFYVE27-dependent manner (By similarity. Regulates centrosome and nuclear positioning during mitotic entry. During the G2 phase of the cell cycle in a BICD2-dependent manner, antagonizes dynein function and drives the separation of nuclei and centrosomes. Required for anterograde axonal transportation of MAPK8IP3/JIP3 which is essential for MAPK8IP3/JIP3 function in axon elongation (By similarity. Through binding with PLEKHM2 and ARL8B, directs lysosome movement toward microtubule plus ends (Probable. Involved in NK cell-mediated cytotoxicity. Drives the polarization of cytolytic granules and microtubule-organizing centers (MTOCs toward the immune synapse between effector NK lymphocytes and target cells.
基因ID	3799
基因名	KIF5B
Swiss	P33176
别名	KIF5B;HEL-S-61;KINH;KNS;KNS1;UKHC

产品验证



Western blot analysis of KIF5B expressed in Mouse brain,Rat brain,HeLa using KIF5B Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn