

TAB3 Rabbit mAb

货号: **AYM29027**

产品信息

| | |
|-------|---|
| 反应 | Human,Mouse |
| 宿主 | Rabbit |
| 克隆性 | Monoclonal |
| 预测反应 | |
| 应用 | WB IF/ICC |
| 推荐浓度 | WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200 |
| 理论分子量 | |
| 实测分子量 | 90kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | HepG2,Mouse spleen,Mouse heart,Rat testis |
| 细胞定位 | cytosol,extracellular exosome,plasma membrane |
| 纯化 | Affinity purification |

抗原信息


| | |
|------|---|
| 抗原信息 | Recombinant fusion protein corresponding to Human TAB3. |
| 序列 | QPSPRNQHSLYTATTPSSSPSRGISSQPKPPFSVNPVYITYTQPTGSPCTSPSPRVIPNPTTVFKITVGRATTENLLNLVDQ EERSAAP |

靶点信息

| | |
|------|--|
| 研究背景 | The product of this gene functions in the NF-kappaB signal transduction pathway. The encoded protein, and the similar and functionally redundant protein MAP3K7IP2/TAB2, forms a ternary complex with the protein kinase MAP3K7/TAK1 and either TRAF2 or TRAF6 in response to stimulation with the pro-inflammatory cytokines TNF or IL-1. Subsequent MAP3K7/TAK1 kinase activity triggers a signaling cascade leading to activation of the NF-kappaB transcription factor. The human genome contains a related pseudogene. Alternatively spliced transcript variants have been described, but their biological validity has not been determined. [provided by RefSeq, Jul 2008] |
|------|--|

| | |
|-------|-----------------|
| 基因ID | 257397 |
| 基因名 | TAB3 |
| Swiss | Q8N5C8 |
| 别名 | MAP3K7IP3; NAP1 |

产品验证



Western blot analysis of TAB3 expressed in Mouse spleen, HepG2, Mouse heart using TAB3 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane . Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn