

— ABLYBIO, Help Your Research



CD30 Rabbit mAb

货号: **AYM29039**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200
理论分子量	120kDa
实测分子量	120kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-549
细胞定位	cytoplasm,extracellular exosome,plasma membrane
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human CD30.
------	---

靶点信息

研究背景	The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor is expressed by activated, but not by resting, T and B cells. TRAF2 and TRAF5 can interact with this receptor, and mediate the signal transduction that leads to the activation of NF-kappaB. This receptor is a positive regulator of apoptosis, and also has been shown to limit the proliferative potential of autoreactive CD8 effector T cells and protect the body against autoimmunity. Two alternatively spliced transcript variants of this gene encoding distinct isoforms have been reported. [provided by RefSeq, Jul 2008]
基因ID	943
基因名	TNFRSF8
Swiss	P28908 (https://www.uniprot.org/uniprotkb/P28908/entry)
别名	CD30,CD30 Rabbit mAb,TNFRSF8,CD30L receptor,Ki-1 antigen,Lymphocyte activation antigen CD30,D1S166E

产品验证



Western blot analysis of CD30 expressed in A-549 using CD30 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)