

— ABLYBIO, Help Your Research



Sterol carrier protein 2 Rabbit mAb

货号: **AYM29044**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	58kDa
实测分子量	58kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,HepG2,U-87MG,Rat liver,Rat brain
细胞定位	Cytoplasm,Mitochondrion,Mitochondrion,Peroxisome
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human Sterol carrier protein 2.
------	---

靶点信息

研究背景	This gene encodes two proteins: sterol carrier protein X (SCPx) and sterol carrier protein 2 (SCP2), as a result of transcription initiation from 2 independently regulated promoters. The transcript initiated from the proximal promoter encodes the longer SCPx protein, and the transcript initiated from the distal promoter encodes the shorter SCP2 protein, with the 2 proteins sharing a common C-terminus. Evidence suggests that the SCPx protein is a peroxisome-associated thiolase that is involved in the oxidation of branched chains in fatty acids, while the SCP2 protein is thought to be an intracellular lipid transfer protein. This gene is highly expressed in organs involved in lipid metabolism, and may play a role in Zellweger syndrome, in which cells are deficient in peroxisomes and have impaired bile acid synthesis. Alternative splicing of this gene produces multiple transcript variants, some encoding different isoforms.
基因ID	6342
基因名	SCP2
Swiss	P22307 (https://www.uniprot.org/uniprotkb/P22307/entry)
别名	Sterol carrier protein 2, Sterol carrier protein 2 Rabbit mAb, SCP2, Acetyl-CoA C-myristoyltransferase, Non-specific lipid-transfer protein, Propanoyl-CoA C-acyltransferase, SCP-2/3-oxoacyl-CoA thiolase, SCP-2/thiolase, SCP-chi, SCPX, Sterol carrier protein X

产品验证



Western blot analysis of Sterol carrier protein 2 expressed in Rat liver, 293T, HepG2 using Sterol carrier protein 2 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/>www.ablybio.cn)