

CD14 Rabbit mAb

货号: **AYM29049**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC ICC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 ICC: 1:50 - 1:200
理论分子量	53kDa
实测分子量	53kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-87MG,THP-1,A-549,RAW264.7
细胞定位	Cell membrane,GPI-anchor,Lipid-anchor,Secreted
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human CD14.
序列	QPHSLDLSHNSLRATVNPSPRCMWSSALNSLNLSFAGLEQVPKGLPAKLRVLDLSCNRLNRPQPDELPEVDNLTLDGN PFLVPGTALPHEGSMNSGVVP

靶点信息


研究背景	The protein encoded by this gene is a surface antigen that is preferentially expressed on monocytes/macrophages. It cooperates with other proteins to mediate the innate immune response to bacterial lipopolysaccharide. Alternative splicing results in multiple transcript variants encoding the same protein. [provided by RefSeq, Mar 2010]
基因ID	929

基因名	CD14
Swiss	P08571
别名	CD14

产品验证



Immunohistochemical analysis of paraffin-embedded human breast, using CD14 Antibody.



Western blot analysis of CD14 expressed in U-87MG, THP-1, A-549 using CD14 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn