

— ABLYBIO, Help Your Research



# TF2B Rabbit mAb

货号: **AYM29056**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IP
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IP:</b> 1:20 - 1:50
理论分子量	34kDa
实测分子量	31kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,Mouse thymus,Rat testis
细胞定位	Nucleus
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein corresponding to Human TF2B.
------	---

## 靶点信息

研究背景	This gene encodes the general transcription factor IIB, one of the ubiquitous factors required for transcription initiation by RNA polymerase II. The protein localizes to the nucleus where it forms a complex (the D AB complex) with transcription factors IID and IIA. Transcription factor IIB serves as a bridge between IID, the factor which initially recognizes the promoter sequence, and RNA polymerase II.
基因ID	2959
基因名	GTF2B
Swiss	Q00403 ( <a href="https://www.uniprot.org/uniprotkb/Q00403/entry">https://www.uniprot.org/uniprotkb/Q00403/entry</a> )
别名	TF2B,TF2B Rabbit mAb,GTF2B,General transcription factor TFIIB,S300-II,TFIIB

## 产品验证



Western blot analysis of TF2B expressed in HeLa, Mouse thymus, Rat testis using TF2B Rabbit mAb at 1:10 00. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/www.ablybio.cn>)