

NR5A2 Rabbit mAb

货号: **AYM29064**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	42kDa/53kDa/56kDa/61kDa
实测分子量	65kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-549,Mouse brain
细胞定位	Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human NR5A2.
序列	SIPHLILELLKCEPDEPQVQAKIMAYLQQEQANRSKHEKLSTFGMLCKMADQTLFSIVEWARSSIFFRELKVDDQMKLLQNC WSELLILDHIYRQVHVGKEGSIFLVTGQQVDYSIIASQAGATLNNLMSHAQELVAKLRSLQFDQREFVCLKFLVLFSLDVKN LENFQLVEGVQEQVNAALLDYTMCNYPQQTEKFGQLLLRLPEIRAISMQAEEYLYKHLNGDVPYNNLLIEMLHAKRA

靶点信息

研究背景	The protein encoded by this gene is a DNA-binding zinc finger transcription factor and is a member of the fushi tarazu factor-1 subfamily of orphan nuclear receptors. The encoded protein is involved in the expression of genes for hepatitis B virus and cholesterol biosynthesis, and may be an important regulator of embryonic development.
基因ID	2494

基因名	NR5A2
Swiss	O00482
别名	NR5A2;B1F;B1F2;CPF;FTF;FTZ-F1;FTZ-F1beta;LRH-1;LRH1;hB1F-2

产品验证



Western blot analysis of NR5A2 expressed in A-549, Mouse brain using NR5A2 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn