

— ABLYBIO, Help Your Research



NONO / p54nrb Rabbit mAb

货号: **AYM29078**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	43kDa/54kDa
实测分子量	54kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	C2C12 cell, mouse skeletal muscle tissue, ROS1728 cells
细胞定位	Nucleus,Nucleus speckle,nucleolus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human NONO / p54nrb.
------	--

靶点信息

研究背景	This gene encodes an RNA-binding protein which plays various roles in the nucleus, including transcriptional regulation and RNA splicing. A rearrangement between this gene and the transcription factor E3 gene has been observed in papillary renal cell carcinoma. Alternatively spliced transcript variants have been described. Pseudogenes exist on Chromosomes 2 and 16.
基因ID	4841
基因名	NONO
Swiss	Q15233 (https://www.uniprot.org/uniprotkb/Q15233/entry)
别名	NONO / p54nrb, NONO / p54nrb Rabbit mAb, NONO, 54 kDa nuclear RNA- and DNA-binding protein, 55 kDa nuclear protein, DNA-binding p52/p100 complex, 52 kDa subunit, NRB54

产品验证



Western blot analysis of NONO / p54nrb expressed in C2C12 cell, mouse skeletal muscle tissue, ROS1728 cells using NONO / p54nrb Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/>www.ablybio.cn)