

# NONO / p54nrb Rabbit mAb

货号: **AYM29078**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:200 <b>IF/ICC:</b> 1:50 - 1:200
理论分子量	43kDa/54kDa
实测分子量	54kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	C2C12 cell, mouse skeletal muscle tissue, ROS1728 cells
细胞定位	Nucleus,Nucleus speckle,nucleolus
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein corresponding to Human NONO / p54nrb.
序列	RQQEEMRR

## 靶点信息

研究背景	This gene encodes an RNA-binding protein which plays various roles in the nucleus, including transcriptional regulation and RNA splicing. A rearrangement between this gene and the transcription factor E3 gene has been observed in papillary renal cell carcinoma. Alternatively spliced transcript variants have been described. Pseudogenes exist on Chromosomes 2 and 16.
基因ID	4841

基因名	NONO
Swiss	Q15233
别名	NONO;MRXS34;NMT55;NRB54;P54;P54NRB;PPP1R114

## 产品验证



Western blot analysis of NONO / p54nrb expressed in C2C12 cell, mouse skeletal muscle tissue, ROS1728 cells using NONO / p54nrb Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)