

Caspase-6 Rabbit mAb

货号: **AYM29082**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC IP
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 IP: 1:20 - 1:50
理论分子量	18,33kDa
实测分子量	11,33kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Jurkat,A-549,MCF7
细胞定位	Cytoplasm
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human Caspase-6.
序列	GIALIFNHERFFWHLTLPERRGTCADRDNLRRFSDLGFVCKCFNDLKAEEELLKIHEVSTVSHADADCFVCFVFLSHGEGN HIYAYDAKIEIQTLTGLFKGDKCHSLVKGPKIFIIQA


靶点信息

研究背景	This gene encodes a member of the cysteine-aspartic acid protease (caspase) family of enzymes. Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes which undergo proteolytic processing at conserved aspartic acid residues to produce two subunits, large and small, that dimerize to form the active enzyme. This protein is processed by caspases 7, 8 and 10, and is thought to function as a downstream enzyme in the caspase activation cascade. Alternative splicing of this gene results in multiple transcript variants that encode different isoforms. [provided by RefSeq, Oct 2015]
基因ID	839
基因名	CASP6
Swiss	P55212
别名	CASP6;MCH2;caspase-6;Caspase-6 p18;Caspase-6

产品验证



Immunohistochemical analysis of paraffin-embedded human colon, using Caspase-6 Antibody.



Western blot analysis of Caspase-6 expressed in Jurkat, A-549, MCF7 using Caspase-6 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn