

GSTO1 Rabbit mAb

货号: **AYM29084**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	23kDa/24kDa/27kDa
实测分子量	28kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	LO2, BxPC-3, HT-29, HeLa, Mouse heart, Rat liver
细胞定位	Cytoplasm, cytosol
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human GSTO1.
序列	MSGESARSLGKGSAPPGPVPEGSIRIYSMRFCFAERTRLVLKAKGIRHEVININLKNKPEWFFKKNPFGLVPVLENSQGQLI YESAITCEYLDEAYPGKLLPDDPYEKACQKMILELFSKVPVSLVGSFIRSQNKEDYAGLKEEFRKEFTKLEEVLTNKKTTFFG GNSISMIDYLIWPWFERLEAMKLNEDVHTPKLKLWMAAMKEDPTVSALLTSEKDWQGFLELYLQNSPEACDYGL

靶点信息

研究背景	The protein encoded by this gene is an omega class glutathione S-transferase (GST) with glutathione-dependent thiol transferase and dehydroascorbate reductase activities. GSTs are involved in the metabolism of xenobiotics and carcinogens. The encoded protein acts as a homodimer and is found in the cytoplasm. Three transcript variants encoding different isoforms have been found for this gene.
基因ID	9446

基因名	GSTO1
Swiss	P78417
别名	GSTO1;GSTO 1-1;GSTTLp28;HEL-S-21;P28;SPG-R;GSTO1-1

产品验证



Western blot analysis of GSTO1 expressed in LO2, BxPC-3, HT-29 using GSTO1 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn