

— ABLYBIO, Help Your Research



# SAFB Rabbit mAb

货号: **AYM29096**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:200 <b>IF/ICC:</b> 1:50 - 1:200
理论分子量	95kDa/102kDa
实测分子量	150kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-87MG,DU145,NIH/3T3,Mouse brain,Mouse lung,Rat brain
细胞定位	Nucleus
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein corresponding to Human SAFB.
------	---------------------------------------------------------

## 靶点信息

研究背景	This gene encodes a DNA-binding protein which has high specificity for scaffold or matrix attachment region on DNA elements (S/MAR DNA). This protein is thought to be involved in attaching the base of chromatin loops to the nuclear matrix but there is conflicting evidence as to whether this protein is a component of chromatin or a nuclear matrix protein. Scaffold attachment factors are a specific subset of nuclear matrix proteins (NMP) that specifically bind to S/MAR. The encoded protein is thought to serve as a molecular base to assemble a 'transcriptosome complex' in the vicinity of actively transcribed genes. It is involved in the regulation of heat shock protein 27 transcription, can act as an estrogen receptor co-repressor and is a candidate for breast tumorigenesis. This gene is arranged head-to-head with a similar gene whose product has the same functions. Multiple transcript variants encoding different isoforms have been found for this gene.
基因ID	6294
基因名	SAFB
Swiss	Q15424 ( <a href="https://www.uniprot.org/uniprotkb/Q15424/entry">https://www.uniprot.org/uniprotkb/Q15424/entry</a> )
别名	SAFB,SAFB Rabbit mAb,HSP27 estrogen response element-TATA box-binding protein,HAP,HET,SAFB1

## 产品验证



Western blot analysis of SAFB expressed in Mouse brain,Rat brain,U-87MG using SAFB Rabbit mAb at 1:10 00. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/www.ablybio.cn>)