

CNTN2 / TAG1 Rabbit mAb

货号: **AYM29111**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IF/ICC FC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200 FC: 1:20 - 1:50
理论分子量	113kDa
实测分子量	135kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse brain,Mouse spinal cord,Rat brain
细胞定位	Cell membrane,GPI-anchor,Lipid-anchor
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human CNTN2 / TAG1.
序列	EVKIRSYNRRRGDGPESLTLVYSAEEEEPRVAPTKVWAKGVSSSEMNVTWEPVQQDMNGILLGYEIRYWKAGDKAAADR VRTAGLDTSARVSGLHPNTKYHVTVRAYNRAGTGPASPSANATMKPPRRPPGNISWTFSSSSLSIKWDPVVPFRNESA VTGYKMLYQNDLHPTLHPTLHGTGKNWIEIPVPEDIGHALVQIRTTGPGGDGIPAEVHIVRNGGTSMVMEN

靶点信息

研究背景	The protein encoded by this gene is a member of the immunoglobulin superfamily. It is a glycosylphosphatidylinositol (GPI)-anchored neuronal membrane protein that functions as a cell adhesion molecule. It may play a role in the formation of axon connections in the developing nervous system. It may also be involved in glial tumorigenesis and may provide a potential target for therapeutic intervention.
------	---

基因ID	6900
基因名	CNTN2
Swiss	Q02246
别名	CNTN2;AXT;FAME5;TAG-1;TAX;TAX1

产品验证



Western blot analysis of CNTN2 / TAG1 expressed in Mouse brain, Mouse spinal cord, Rat brain using CNTN2 / TAG1 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn