

— ABLYBIO, Help Your Research



ZA2G / AZGP1 Rabbit mAb

货号: **AYM29118**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IP
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IP: 1:20 - 1:50
理论分子量	34kDa
实测分子量	43kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	Human serum
细胞定位	Secreted
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human ZA2G / AZGP1.
------	---

靶点信息

研究背景	Zinc- α -2-Glycoprotein (AZGP1) can be found in blood plasma, seminal plasma, urine, sweat, saliva, liver, and epithelial cells of various human glands. AZGP1 has been proposed in the regulation of body weight, and the melanin production by normal and malignant melanocytes. AZGP1 stimulates lipid degradation in adipocytes and causes the extensive fat losses associated with some advanced cancers. AZGP1 has been reported to stimulate lipid breakdown and may have an important role in lipid homeostasis. Mature human AZGP1 consists of one MHC class I antigen region and a C2-type Ig-like domain. AZGP1 has two alternative splice forms, one shows a 66 amino acids substitution for the C-terminal 30 amino acids, the other one shows a nine Lys substitution for amino acid 151-298.
基因ID	563
基因名	AZGP1
Swiss	P25311
别名	ZA2G / AZGP1,ZA2G / AZGP1 Rabbit mAb,AZGP1,ZAG,ZNGP1

产品验证



Western blot analysis of ZA2G / AZGP1 expressed in Human serum using ZA2G / AZGP1 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/>www.ablybio.cn)