

— ABLYBIO, Help Your Research



TIP49A Rabbit mAb

货号: **AYM29134**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB FC
推荐浓度	WB: 1:500 - 1:2000 FC: 1:20 - 1:50
理论分子量	42kDa/50kDa
实测分子量	50kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HL-60,MCF7,HepG2,293T,Mouse spleen,Mouse skeletal muscle,Mouse thymus,Mouse brain
细胞定位	Cytoplasm,Membrane,Nucleus,Nucleus matrix,centrosome,cytoskeleton,microtubule organizing center,nucleoplasm
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human TIP49A.
------	---

靶点信息

研究背景	This gene encodes a protein that has both DNA-dependent ATPase and DNA helicase activities and belongs to the ATPases associated with diverse cellular activities (AAA+) protein family. The encoded protein associates with several multisubunit transcriptional complexes and with protein complexes involved in both ATP-dependent remodeling and histone modification. Alternate splicing results in multiple transcript variants.
基因ID	8607
基因名	RUVBL1
Swiss	Q9Y265 (https://www.uniprot.org/uniprotkb/Q9Y265/entry)
别名	TIP49A,TIP49A Rabbit mAb,RUVBL1,49 kDa TATA box-binding protein-interacting protein,54 kDa erythrocyte cytosolic protein,INO80 complex subunit H,Nuclear matrix protein 238,Pontin 52,TIP60-associated protein 54-alpha,INO80H,NMP238,TIP49

产品验证



Western blot analysis of TIP49A expressed in Mouse spleen,HL-60,MCF7 using TIP49A Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/>www.ablybio.cn)