

TIP49A Rabbit mAb

货号: **AYM29134**

产品信息

| | |
|-------|---|
| 反应 | Human,Mouse,Rat |
| 宿主 | Rabbit |
| 克隆性 | Monoclonal |
| 预测反应 | |
| 应用 | WB FC |
| 推荐浓度 | WB: 1:500 - 1:2000 FC: 1:20 - 1:50 |
| 理论分子量 | 42kDa/50kDa |
| 实测分子量 | 50kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | HL-60,MCF7,HepG2,293T,Mouse spleen,Mouse skeletal muscle,Mouse thymus,Mouse brain |
| 细胞定位 | Cytoplasm,Membrane,Nucleus,Nucleus matrix,centrosome,cytoskeleton,microtubule organizing center,nucleoplasm |
| 纯化 | Affinity purification |

抗原信息


| | |
|------|---|
| 抗原信息 | Recombinant fusion protein corresponding to Human TIP49A. |
|------|---|

靶点信息

| | |
|------|--|
| 研究背景 | This gene encodes a protein that has both DNA-dependent ATPase and DNA helicase activities and belongs to the ATPases associated with diverse cellular activities (AAA+) protein family. The encoded protein associates with several multisubunit transcriptional complexes and with protein complexes involved in both ATP-dependent remodeling and histone modification. Alternate splicing results in multiple transcript variants. |
| 基因ID | 8607 |
| 基因名 | RUVBL1 |

| | |
|-------|--------|
| Swiss | Q9Y265 |
| 别名 | TIP49A |

产品验证



Western blot analysis of TIP49A expressed in Mouse spleen,HL-60,MCF7 using TIP49A Rabbit mAb at 1:1 000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. B locking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn