

SPINK5 / LEKTI Rabbit mAb

货号: **AYM29160**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	104kDa/120kDa/124kDa
实测分子量	121kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,MCF7,RAW264.7,Mouse skin,Mouse stomach,Rat stomach
细胞定位	Secreted
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human SPINK5 / LEKTI.
序列	FKKGERDGDGFICPDYYEAVCGTDGKTYDNRCALCAENAKTGSQIGVKSEGECKSSNPEQDVCSAFRPFVRDGRGLGCTRE NDPVLGPDGKTHGNKCAMCAEL

靶点信息

研究背景	This gene encodes a multidomain serine protease inhibitor that contains 15 potential inhibitory domains. The encoded preproprotein is proteolytically processed to generate multiple protein products, which may exhibit unique activities and specificities. These proteins may play a role in skin and hair morphogenesis, as well as anti-inflammatory and antimicrobial protection of mucous epithelia. Mutations in this gene may result in Netherton syndrome, a disorder characterized by ichthyosis, defective cornification, and atopy. This gene is present in a gene cluster on chromosome 5. Alternative splicing results in multiple transcript variants.
------	--

基因ID	11005
基因名	SPINK5
Swiss	Q9NQ38
别名	SPINK5;LEKTI;LETKI;NETS;NS;VAKTI

产品验证



Western blot analysis of SPINK5 / LEKTI expressed in Mouse skin,Rat stomach,HeLa using SPINK5 / LEKTI Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn