

Pumilio 2 Rabbit mAb

货号: **AYM29166**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IF/ICC IP FC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200 IP: 1:20 - 1:50 FC: 1:20 - 1:50
理论分子量	114kDa
实测分子量	114kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	PC-3,Mouse testis,Rat testis
细胞定位	cytoplasm,cytoplasmic stress granule,cytosol,nuclear membrane,perinuclear region of cytoplasm
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human Pumilio 2.
序列	MNHDFQALALESRGMGELLPTKKFWEPDDSTKDGQKGI FLGDDEWRETAWGASHHMSQPIMVQRRSGQGFHGNSE VNAILSPRESGGLGVSMVEYVLS

靶点信息

研究背景	This gene encodes a protein that belongs to a family of RNA-binding proteins. The encoded protein functions as a translational repressor during embryonic development and cell differentiation. This protein is also thought to be a positive regulator of cell proliferation in adipose-derived stem cells. Alternate splicing results in multiple transcript variants. [provided by RefSeq, Sep 2013]
------	---

基因ID	23369
基因名	PUM2
Swiss	Q8TB72
别名	PUMH2; PUML2

产品验证



Western blot analysis of Pumilio 2 expressed in PC-3, Mouse testis, Rat testis using Pumilio 2 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120 S.

实验步骤

访问官网浏览详情: www.ablybio.cn