

RNF14 Rabbit mAb

货号: **AYM29168**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	
实测分子量	54kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	SH-SY5Y,Jurkat,K-562,Mouse placenta,Rat brain
细胞定位	cytoplasm,cytosol,nucleoplasm,nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human RNF14.
序列	LDLMADVVCPRPCCQLPVMQEPGCTMGICSSCNFAFCTLCRLTYHGVSPCKVTAEKLMDLRNEYLQADEANKRLLDQRY GKRVIQKALEEMESKEWLEKN

靶点信息

研究背景	The protein encoded by this gene contains a RING zinc finger, a motif known to be involved in protein-protein interactions. This protein interacts with androgen receptor (AR) and may function as a coactivator that induces AR target gene expression in prostate. A dominant negative mutant of this gene has been demonstrated to inhibit the AR-mediated growth of prostate cancer. This protein also interacts with class III ubiquitin-conjugating enzymes (E2s) and may act as a ubiquitin-ligase (E3) in the ubiquitination of certain nuclear proteins. Six alternatively spliced transcript variants encoding two distinct isoforms have been reported. [provided by RefSeq, Jan 2011]
------	---

基因ID	9604
基因名	RNF14
Swiss	Q9UBS8
别名	RNF14;ARA54;HFB30;HRIHFB2038;TRIAD2

产品验证



Western blot analysis of RNF14 expressed in Mouse placenta,Rat brain,SH-SY5Y using RNF14 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn