

Pancreatic Lipase (PNLIP) Rabbit mAb

货号: **AYM29179**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200
理论分子量	51kDa
实测分子量	49kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	BxPC-3
细胞定位	Secreted
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human Pancreatic Lipase (PNLIP) Rabbit mAb.
序列	IGHSLGAHAAGEAGRRRTNGTIGRITGLDPAEPCFQGTPELVRLDPSDAKFVDVIHTDGAPIVPNLFGMSQVVGHLDFFPN GGVEMPGCKKNILSQIVDIDGIWEGTRDFAACNHLRSYKYTDSIVNPDGFAGFPCASYNVFTANKCFPCPSGGCPQMGH YADRYPGKTNDVGQKFYLDTGASNFAWRVKVSVTLSGKKVTGHILVSLFGNKGNSKQYEIFKGTLPDSTHSNEFDS DVDVGLDQMVKFIWYNNVINPTLPRVGASKIIVETNVGKQFNFCSPETVREEVLLTLTPC

靶点信息

研究背景	This gene is a member of the lipase gene family. It encodes a carboxyl esterase that hydrolyzes insoluble , emulsified triglycerides, and is essential for the efficient digestion of dietary fats. This gene is expressed specifically in the pancreas.
基因ID	5406

基因名	PNLIP
Swiss	P16233
别名	PNLIP;PL;PNLIPD;PTL

产品验证



Western blot analysis of Pancreatic Lipase (PNLIP) Rabbit expressed in BxPC-3 using Pancreatic Lipase (PNLIP) Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn