

IKZF3 Rabbit mAb

货号: **AYM29184**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200
理论分子量	29-58kDa
实测分子量	65kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	BT-474,Mouse thymus
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human IKZF3.
序列	HCFDVNYNSSYMYEKESELIQTRMMDQAINNAISYLGAEALRPLVQTPAPTSEMVPVISSMYPIALTRAEMSNGAPQELEK KSIHLPEKSVPSERGLSPNNSGHDSTDTDSNHEERQNHIIYQQNHMVLSRARNGMPLLKEVPRSYELLKPPPICPRDSVKVI NKEGEVMD

靶点信息

研究背景	This gene encodes a member of the Ikaros family of zinc-finger proteins. Three members of this protein family (Ikaros, Aiolos and Helios) are hematopoietic-specific transcription factors involved in the regulation of lymphocyte development. This gene product is a transcription factor that is important in the regulation of B lymphocyte proliferation and differentiation. Both Ikaros and Aiolos can participate in chromatin remodeling. Regulation of gene expression in B lymphocytes by Aiolos is complex as it appears to require the sequential formation of Ikaros homodimers, Ikaros/Aiolos heterodimers, and Aiolos homodimers. Several alternative transcripts encoding different isoforms have been described, as well as some non-protein coding variants.
基因ID	22806
基因名	IKZF3
Swiss	Q9UKT9
别名	IKZF3;AIO;AIOLOS;ZNFN1A3

产品验证



Western blot analysis of IKZF3 expressed in BT-474, Mouse thymus using IKZF3 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn