

— ABLYBIO, Help Your Research



PSMD14 Rabbit mAb

货号: **AYM29189**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	35kDa
实测分子量	35kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,Mouse brain,Mouse thymus,Rat lung,Rat brain
细胞定位	cytosol,cytosolic proteasome complex,extracellular region,nucleoplasm,nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human PSMD14.
------	---

靶点信息

研究背景	This gene encodes a component of the 26S proteasome. The 26S proteasome is a large multiprotein complex that catalyzes the degradation of ubiquitinated intracellular proteins. The encoded protein is a component of the 19S regulatory cap complex of the 26S proteasome and mediates substrate deubiquitination. A pseudogene of this gene is also located on the long arm of chromosome 2. [provided by RefSeq, Feb 2012]
基因ID	10213
基因名	PSMD14
Swiss	O00487 (https://www.uniprot.org/uniprotkb/O00487/entry)
别名	PSMD14,PSMD14 Rabbit mAb,26S proteasome non-ATPase regulatory subunit 14,26S proteasome regulatory subunit RPN11,26S proteasome-associated PAD1 homolog 1,POH1

产品验证



Western blot analysis of PSMD14 expressed in Mouse brain,Rat lung,HeLa using PSMD14 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)