

— ABLYBIO, Help Your Research



TAF15 Rabbit mAb

货号: **AYM29197**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 FC: 1:20 - 1:50
理论分子量	61kDa
实测分子量	75kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	SKOV3,Jurkat
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human TAF15.
------	--

靶点信息

研究背景	This gene encodes a member of the TET family of RNA-binding proteins. The encoded protein plays a role in RNA polymerase II gene transcription as a component of a distinct subset of multi-subunit transcription initiation factor TFIID complexes. Translocations involving this gene play a role in acute leukemia and extr askeletal myxoid chondrosarcoma, and mutations in this gene may play a role in amyotrophic lateral scler osis. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gen e.
基因ID	8148
基因名	TAF15
Swiss	Q92804 (https://www.uniprot.org/uniprotkb/Q92804/entry)
别名	TAF15,TAF15 Rabbit mAb,68 kDa TATA-binding protein-associated factor,RNA-binding protein 56,RBP56,TA F2N

产品验证



Western blot analysis of TAF15 expressed in SKOV3, Jurkat using TAF15 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/>www.ablybio.cn)