

— ABLYBIO, Help Your Research



# PDCD7 Rabbit mAb

货号: **AYM29228**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:200 <b>IF/ICC:</b> 1:50 - 1:200
理论分子量	55kDa
实测分子量	55kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	TE-1,SKOV3,Mouse testis,Mouse thymus,Rat brain
细胞定位	Nucleus
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein corresponding to Human PDCD7.
------	--

## 靶点信息

研究背景	This gene encodes a 59 kDa protein that is associated with the U11 small nuclear ribonucleoprotein (snRNP), which is a component of the minor U12-type spliceosome responsible for catalyzing pre-mRNA splicing of U12-type introns.
基因ID	10081
基因名	PDCD7
Swiss	Q8N8D1 ( <a href="https://www.uniprot.org/uniprotkb/Q8N8D1/entry">https://www.uniprot.org/uniprotkb/Q8N8D1/entry</a> )
别名	PDCD7,PDCD7 Rabbit mAb,ES18

## 产品验证



Western blot analysis of PDCD7 expressed in Mouse testis,Rat brain,TE-1 using PDCD7 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/>[www.ablybio.cn](http://www.ablybio.cn))