

PDCD7 Rabbit mAb

货号: **AYM29228**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	55kDa
实测分子量	55kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	TE-1,SKOV3,Mouse testis,Mouse thymus,Rat brain
细胞定位	Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human PDCD7.
序列	MALPPFFGQGRPGPPPPQPPPPAPFGCPPPLPSAFPPPLPQRPGPFPGASAPFLQPPLALQPRASAEASRGGGGAGAFY PVPPPLPPPPQCRPFG

靶点信息

研究背景	This gene encodes a 59 kDa protein that is associated with the U11 small nuclear ribonucleoprotein (snRNP), which is a component of the minor U12-type spliceosome responsible for catalyzing pre-mRNA splicing of U12-type introns.
基因ID	10081

基因名	PDCD7
Swiss	Q8N8D1
别名	PDCD7; ES18; HES18; programmed cell death 7

产品验证



Western blot analysis of PDCD7 expressed in Mouse testis, Rat brain, TE-1 using PDCD7 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn