

NSDHL Rabbit mAb

货号: **AYM29231**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	41kDa
实测分子量	38kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T
细胞定位	endoplasmic reticulum,lipid droplet
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human NSDHL.
序列	MEPAVSEPMRDQVARTHLTEDTPKVNADIEKVNQNQAKRCTVIGGSGFLGQHMVEQLLARGYAVNVFDIQGGFDNPQV RFFLGDLCSRQDLYPALKGVNTVVFHCASPPSSNNKELFYRVNYIGTKNVIETCKEAGVQKLILTSSASVIFEGVDIKNGTED LPYAMKPIDYYTETKILQERAVLGANDPEKNFLTТАIRPHGIFGPRDPQLVPILIEAARNGMKMFVINGKNLVDFTFV

靶点信息

研究背景	The protein encoded by this gene is localized in the endoplasmic reticulum and is involved in cholesterol biosynthesis. Mutations in this gene are associated with CHILD syndrome, which is a X-linked dominant disorder of lipid metabolism with disturbed cholesterol biosynthesis, and typically lethal in males. Alternatively spliced transcript variants with differing 5' UTR have been found for this gene.
------	---

基因ID	50814
基因名	NSDHL
Swiss	Q15738
别名	NSDHL;H105E3;SDR31E1;XAP104

产品验证



Western blot analysis of NSDHL expressed in 293T using NSDHL Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn