

CD79b Rabbit mAb

货号: **AYM29268**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IF/ICC FC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200 FC: 1:20 - 1:50
理论分子量	14kDa/26kDa
实测分子量	30-50kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Raji
细胞定位	Cell membrane,Single-pass type I membrane protein
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human CD79b.
序列	SEVYQCGGTGLRVMGFSTLAQLKQRNTLKDGIIMIQTLLIILFIIVPIFLLLDKDDSKAGMEEDHTYEGLDIDQTATYEDIVTLR TGEVKWSVGEHPGQE

靶点信息

研究背景	The B lymphocyte antigen receptor is a multimeric complex that includes the antigen-specific component , surface immunoglobulin (Ig). Surface Ig non-covalently associates with two other proteins, Ig-alpha and I g-beta, which are necessary for expression and function of the B-cell antigen receptor. This gene encodes the Ig-beta protein of the B-cell antigen component. Alternatively spliced transcript variants encoding different isoforms have been described.
------	--

基因ID	974
基因名	CD79B
Swiss	P40259
别名	CD79B;AGM6;B29;IGB

产品验证



Western blot analysis of CD79b expressed in Raji using CD79b Rabbit mAb at 1:1000. Secondary antibody : HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn