

IFNGR1 Rabbit mAb

货号: **AYM29271**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 FC: 1:20 - 1:50
理论分子量	21kDa/54kDa
实测分子量	45-100kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,Mouse liver,Rat liver
细胞定位	Membrane,Single-pass type I membrane protein
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human IFNGR1.
序列	IKKINPLKEKSIILPKSLISVVRSATLETKPESKYVSLITSYQPFSLKEKVVCEEPLSPATVPGMHTEDNPGKVEHTEELSSITEVV TTEENIPDVVPGSHLTPIERESSPLSSNQSEPGSIALNSYHSRNCSESDHSRNGFDTDSSCLESHTSLDSEFPNPKGEI KTEGQELITVIKAPTSFGYDKPHVLVDLLVDDSGKESLIGYRPTEDSKEFS

靶点信息

研究背景	This gene (IFNGR1) encodes the ligand-binding chain (alpha) of the gamma interferon receptor. Human interferon-gamma receptor is a heterodimer of IFNGR1 and IFNGR2. A genetic variation in IFNGR1 is associated with susceptibility to Helicobacter pylori infection. In addition, defects in IFNGR1 are a cause of Mendelian susceptibility to mycobacterial disease, also known as familial disseminated atypical mycobacterial infection.
------	---

基因ID	3459
基因名	IFNGR1
Swiss	P15260
别名	IFNGR1;CD119;IFNGR;IMD27A;IMD27B

产品验证



Western blot analysis of IFNGR1 expressed in 293T, Mouse liver, Rat liver using IFNGR1 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn