

RIP2 Rabbit mAb

货号: **AYM29274**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	45kDa/61kDa
实测分子量	61kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse lung,Mouse brain,Rat brain
细胞定位	Cytoplasm
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human RIP2.
序列	MNGEAICSALEPTIPYHKLADLRYLSRGASGTVSSARHADWRVQVAVKHLHIHTPLLDSEKDVLRREAEILHKARFSYILPILGI CNEPEFLGIVTEYMPNGSLNELLHRKTEYPDVAWPLRFRILHEIALGVNYLHNMTPELLHDLKTQNILLDNEFHVKIADFGL SKWRMMSLSQSRSSKSAPEGGTIIYMPPENYEPGQKSRASIKHDIYSYAVITWEVLSRKQPFEDVTNPLQIMYSVSQGHRP VINEESLPYDIPHRARMISLIESGWAQNPDER

靶点信息

研究背景	This gene encodes a member of the receptor-interacting protein (RIP) family of serine/threonine protein kinases. The encoded protein contains a C-terminal caspase activation and recruitment domain (CARD), and is a component of signaling complexes in both the innate and adaptive immune pathways. It is a potent activator of NF-kappaB and inducer of apoptosis in response to various stimuli.
基因ID	8767

基因名	RIPK2
Swiss	O43353
别名	RIPK2;CARD3;CARDIAK;CCK;GIG30;RICK;RIP2

产品验证



Western blot analysis of RIP2 expressed in Mouse lung, Mouse brain, Rat brain using RIP2 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn