

USF1 Rabbit mAb

货号: **AYM29286**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	27kDa/33kDa
实测分子量	34kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,NIH/3T3,Jurkat,K-562,THP-1,Rat spleen
细胞定位	Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human USF1.
序列	TSDDAVDTEGTAETHYTFPSTAVGDGAGGTTSGSTAAVTTQGSEALLGQATPPGTGQFFVMMSPQEV LQGGSQRS IAPRTHPYSPKSEAPRTRDEKR

靶点信息

研究背景	This gene encodes a member of the basic helix-loop-helix leucine zipper family, and can function as a cellular transcription factor. The encoded protein can activate transcription through pyrimidine-rich initiator (Inr) elements and E-box motifs. This gene has been linked to familial combined hyperlipidemia (FCHL). Alternative splicing of this gene results in multiple transcript variants. A related pseudogene has been defined on chromosome 21.
基因ID	7391

基因名	USF1
Swiss	P22415
别名	USF1;FCHL;FCHL1;HYPLIP1;MLTF;MLTFI;UEF;bHLHb11

产品验证



Western blot analysis of USF1 expressed in Rat spleen, HeLa, NIH/3T3 using USF1 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn