

FADD Rabbit mAb

货号: **AYM29289**

产品信息

反应	Mouse
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200
理论分子量	28kDa
实测分子量	28kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	NIH/3T3,Mouse brain,Mouse thymus
细胞定位	cytoplasm,cytosol,neuron projection,nucleus,plasma membrane,riposome
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Mouse FADD .
序列	MDPFLVLLHLSLGSLSGNDLMEKFLCRERVSKRKLERVQSLDLFTVLLQNDLERGHTGLLRELLASLRRHDLQLRLLD FEAGTATAAPPGEADLQVAFDIVCDNVGRDWKRLARELKVSEAKMDGIEEKYPRLSERVRESLKVWKNAEKKNASVAGL VKALRTCLNVLVADLVVEEAQESVSKSENMSPVLRDSTVSSSETP

靶点信息

研究背景	Apoptotic adaptor molecule that recruits caspase-8 or caspase-10 to the activated Fas (CD95 or TNFR-1) receptors. The resulting aggregate called the death-inducing signaling complex (DISC) performs caspase-8 proteolytic activation. Active caspase-8 initiates the subsequent cascade of caspases mediating apoptosis. Involved in interferon-mediated antiviral immune response, playing a role in the positive regulation of interferon signaling.
------	----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

基因ID	14082
基因名	
Swiss	Q61160
别名	Mort1/FADD

产品验证



Western blot analysis of FADD expressed in NIH/3T3, Mouse brain, Mouse thymus using FADD Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn