

# MST1 Rabbit mAb

货号: **AYM29297**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC FC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:200 <b>FC:</b> 1:20 - 1:50
理论分子量	80kDa
实测分子量	85kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse kidney,Rat kidney
细胞定位	Secreted
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein corresponding to Human MST1.
序列	CREAACVWCNGEERYGAVDRTESGRECQRWDLQHPHQHPFEPGKFLDQGLDDNYCRNPDGSERPWCYTTDPQIEREF CDLPRCGSEAQPRQEATTVSCFRGKGEYRGRTANTTTAGVPCQRWDAQIPHQHRFTPEKYACKDLRENFCRNPDGSEAP WCFTLRPGMRAAFYQIRRCTDDVVRPQDCYHGAGEQYRGTVSKTRKGVQCQRWSAETPHKPFQFTTSEPHAQLEENFC RNPDGDSHGPPWCYTMDPRTPFDYCALRRCADQPPSILDPPDQVQFEKCGKRVDRLDQRRSKLRVVG

## 靶点信息

研究背景	The protein encoded by this gene contains four kringle domains and a serine protease domain, similar to that found in hepatic growth factor. Despite the presence of the serine protease domain, the encoded protein may not have any proteolytic activity. The receptor for this protein is RON tyrosine kinase, which upon activation stimulates ciliary motility of ciliated epithelial lung cells. This protein is secreted and cleaved to form an alpha chain and a beta chain bridged by disulfide bonds.
------	---

基因ID	4485
基因名	MST1
Swiss	P26927
别名	MST1;D3F15S2;DNF15S2;HGFL;MSP;NF15S2

### 产品验证



Western blot analysis of MST1 expressed in Mouse kidney,Rat kidney using MST1 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

### 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)