

HSF2 Rabbit mAb

货号: **AYM29336**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IF/ICC FC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200 FC: 1:20 - 1:50
理论分子量	58kDa/60kDa
实测分子量	60kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	NIH/3T3, RAW264.7, Rat testis
细胞定位	Cytoplasm, Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human HSF2.
序列	LDEQRFAKEILPKYFKHNNMASFVRQLNMYGFRKVVHIDSGIVKQERDGPVEFQHPYFKQGQDDLLENIKRKVSSSKPEE NKIRQEDLTKIISSAQKVQIKQETIESRSLSELKSENESESLWKEVSELRAKHAQQQVIRKIVQFIVTLVQNNQLVSLKRKRPLLL NTNGAQKKNLFQH

靶点信息

研究背景	The protein encoded by this gene belongs to the HSF family of transcription factors that bind specifically to the heat-shock promoter element and activate transcription. Heat shock transcription factors activate heat-shock response genes under conditions of heat or other stresses. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.
------	--

基因ID	3298
基因名	HSF2
Swiss	Q03933
别名	HSF2;HSF 2;HSTF 2;HSTF2

产品验证



Western blot analysis of HSF2 expressed in NIH/3T3,RAW264.7,Rat testis using HSF2 Rabbit mAb at 1:1 000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. B locking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn