

— ABLYBIO, Help Your Research



# ReIB Rabbit mAb

货号: **AYM29349**

## 产品信息

|       |  |
|-------|--|
| 反应    | Human  |
| 宿主    | Rabbit   |
| 克隆性   | Monoclonal   |
| 预测反应  |  |
| 应用    | WB IHC IF/ICC IP   |
| 推荐浓度  | <b>WB:</b> 1:500 - 1:2000<br><b>IHC:</b> 1:50 - 1:200<br><b>IF/ICC:</b> 1:50 - 1:200<br><b>IP:</b> 1:20 - 1:50 |
| 理论分子量 | 62kDa  |
| 实测分子量 | 70kDa  |
| 形式    | Liquid   |
| 保存条件  | Store at -20°C. Avoid freeze / thaw cycles.<br>Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.                  |
| 偶联物   | Unconjugated   |
| 阳性对照  | 293F,Mouse thymus  |
| 细胞定位  | Cytoplasm,Nucleus,centrosome,cytoskeleton,microtubule organizing center  |
| 纯化    | Affinity purification  |


## 抗原信息

|      |   |
|------|---|
| 抗原信息 | Recombinant fusion protein corresponding to Human ReIB. |
|------|---|


## 靶点信息

|       |  |
|-------|--|
| 研究背景  | NF-kappa-B is a pleiotropic transcription factor which is present in almost all cell types and is involved in many biological processes such as inflammation, immunity, differentiation, cell growth, tumorigenesis and apoptosis. NF-kappa-B is a homo- or heterodimeric complex formed by the Rel-like domain-containing proteins RELA/p65, RELB, NFKB1/p105, NFKB1/p50, REL and NFKB2/p52. NF-kappa-B is controlled by various mechanisms of post-translational modification and subcellular compartmentalization as well as by interactions with other cofactors or corepressors. NF-kappa-B complexes are held in the cytoplasm in an inactive state complexed with members of the NF-kappa-B inhibitor (I-kappa-B) family. In a conventional activation pathway, I-kappa-B is phosphorylated by I-kappa-B kinases (IKKs) in response to different activators, subsequently degraded thus liberating the active NF-kappa-B complex which translocates to the nucleus. NF-kappa-B heterodimeric RelB-p50 and RelB-p52 complexes are transcriptional activators. RELB neither associates with DNA nor with RELA/p65 or REL. Stimulates promoter activity in the presence of NFKB2/p49. Increased repression of the heterodimer is seen in the presence of NFKB2/p52. Is required for both T and B lymphocyte maturation and function. |
| 基因ID  | 5971   |
| 基因名   | RELB   |
| Swiss | Q01201 ( <a href="https://www.uniprot.org/uniprotkb/Q01201/entry">https://www.uniprot.org/uniprotkb/Q01201/entry</a> )   |
| 别名    | RelB,RelB Rabbit mAb,I-Rel   |

## 产品验证



Western blot analysis of RelB expressed in 293F, Mouse thymus using RelB Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.



Immunohistochemical analysis of paraffin-embedded human tonsil, using RelB Antibody.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/www.ablybio.cn>)