

# ReIB Rabbit mAb

货号: **AYM29349**

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC IP
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:200 <b>IF/ICC:</b> 1:50 - 1:200 <b>IP:</b> 1:20 - 1:50
理论分子量	62kDa
实测分子量	70kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	293F, Mouse thymus
细胞定位	Cytoplasm, Nucleus, centrosome, cytoskeleton, microtubule organizing center
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein corresponding to Human ReIB.
序列	GPGPQPHLVITEQPKQRGMRFRYECEGRSAGSILGESSTEASKTLPALIELRDCGGLREVEVTA CLVWKDWPHRVHPHSLV GKDCTDGCICRVRLRPHVSPRHSFNNGIQCVRKKEIEAAIERKIQLGIDPYNAGSLKNHQEVDMMNVVVICFQASYRDQQGQ MRRMDPVLSEPVYDKKSTNTSELRICRINKESGPCTGGEELYLLCDKQVKEDISVVFSTRASWEGRADFSQADVHRQIAIVF KTPPYEDLEIVEPVTNNVFLQRLTDGVCSEPLPFTYLPR


## 靶点信息

研究背景	NF-kappa-B is a pleiotropic transcription factor which is present in almost all cell types and is involved in many biological processes such as inflammation, immunity, differentiation, cell growth, tumorigenesis and apoptosis. NF-kappa-B is a homo- or heterodimeric complex formed by the Rel-like domain-containing proteins RELA/p65, RELB, NFKB1/p105, NFKB1/p50, REL and NFKB2/p52. NF-kappa-B is controlled by various mechanisms of post-translational modification and subcellular compartmentalization as well as by interactions with other cofactors or corepressors. NF-kappa-B complexes are held in the cytoplasm in an inactive state complexed with members of the NF-kappa-B inhibitor (I-kappa-B) family. In a conventional activation pathway, I-kappa-B is phosphorylated by I-kappa-B kinases (IKKs) in response to different activators, subsequently degraded thus liberating the active NF-kappa-B complex which translocates to the nucleus. NF-kappa-B heterodimeric RelB-p50 and RelB-p52 complexes are transcriptional activators. RELB neither associates with DNA nor with RELA/p65 or REL. Stimulates promoter activity in the presence of NFKB2/p49. Increased repression of the heterodimer is seen in the presence of NFKB2/p52. Is required for both T and B lymphocyte maturation and function.
基因ID	5971
基因名	RELB
Swiss	Q01201
别名	RELB;I-REL;IREL;REL-B

## 产品验证



Immunohistochemical analysis of paraffin-embedded human tonsil, using RelB Antibody.



Western blot analysis of RelB expressed in 293F, Mouse thymus using RelB Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)