

CREB Rabbit mAb

货号: **AYM29354**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC IP FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 IP: 1:20 - 1:50 FC: 1:20 - 1:50
理论分子量	25kDa/35kDa/36kDa
实测分子量	40kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-431
细胞定位	Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human CREB.
序列	MTMESGAENQQSGDAAVTEAENQQMTVQAQPQIATLAQVSMPPAAHATSSAPTPTLVQLPNGQTVQVHGVIQAAQPSVI QSPQVQTVQSSCKDLKRLFSGTQISTIAESEDSESVDVSD

靶点信息


研究背景	This gene encodes a transcription factor that is a member of the leucine zipper family of DNA binding proteins. This protein binds as a homodimer to the cAMP-responsive element, an octameric palindrome. The protein is phosphorylated by several protein kinases, and induces transcription of genes in response to hormonal stimulation of the cAMP pathway. Alternate splicing of this gene results in several transcript variants encoding different isoforms.
------	--

基因ID	1385
基因名	CREB1
Swiss	P16220
别名	CREB1;CREB;CREB-1

产品验证



Immunohistochemical analysis of paraffin-embedded mouse brain, using CREB Antibody.



Western blot analysis of CREB expressed in A-431 using CREB Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn