

Periostin Rabbit mAb

货号: **AYM29362**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB FC
推荐浓度	WB: 1:500 - 1:2000 FC: 1:20 - 1:50
理论分子量	80kDa/83kDa/87kDa/90kDa/93kDa
实测分子量	85kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Rat lung
细胞定位	Golgi apparatus,Secreted,extracellular matrix,extracellular space
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human Periostin.
序列	CMEKGSKQGRNGAIHIFREIIPAEKSLHEKQDKRFSTFLSLEAADLKELLTQPGDWTLFVPTNDAFKGMTSEEKEILIR DKNALQNIILYHLTPGVFIGKGFEPGVNLTQSGKIFLKEVNDLLVNLKSKESDIMTTNGVIHVVDKLLYPADTPVGN DQLEILNKLKIKYIKFVIRGTFKEIPVTVYKPIIKKYTKIIDGVPVEITEKETREERIITGPEIKYTRISTGGGETEETLKLLQEDT PVRKLQANKKVQGSRRRLREGRSQ

靶点信息

研究背景	This gene encodes a secreted extracellular matrix protein that functions in tissue development and regeneration, including wound healing, and ventricular remodeling following myocardial infarction. The encoded protein binds to integrins to support adhesion and migration of epithelial cells. This protein plays a role in cancer stem cell maintenance and metastasis. Mice lacking this gene exhibit cardiac valve disease, and skeletal and dental defects. Alternative splicing results in multiple transcript variants encoding different isoforms.
------	--

基因ID	10631
基因名	POSTN
Swiss	Q15063
别名	POSTN;OSF-2;OSF2;PDLPOSTN;PN;periostin

产品验证



Western blot analysis of Periostin expressed in Rat lung using Periostin Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn