

ERp57 Rabbit mAb

货号: **AYM29364**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IP
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IP: 1:20 - 1:50
理论分子量	56kDa
实测分子量	57kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	NIH/3T3,MCF7,Jurkat,HepG2,LO2,Mouse brain,Mouse thymus,Mouse liver,Rat liver
细胞定位	Endoplasmic reticulum,Endoplasmic reticulum lumen,Melanosome
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human ERp57.
------	--

靶点信息

研究背景	This gene encodes a protein of the endoplasmic reticulum that interacts with lectin chaperones calreticulin and calnexin to modulate folding of newly synthesized glycoproteins. The protein was once thought to be a phospholipase; however, it has been demonstrated that the protein actually has protein disulfide isomerase activity. It is thought that complexes of lectins and this protein mediate protein folding by promoting formation of disulfide bonds in their glycoprotein substrates. This protein also functions as a molecular chaperone that prevents the formation of protein aggregates.
基因ID	2923

基因名	PDIA3
Swiss	P30101
别名	ERp57

产品验证



Western blot analysis of ERp57 expressed in NIH/3T3, MCF7, Jurkat using ERp57 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn