

A2M Rabbit mAb

货号: **AYM29372**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200
理论分子量	163kDa
实测分子量	163kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse uterus,Mouse brain
细胞定位	Secreted
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human A2M.
序列	ALAGNQDKRKEVLKSLNEEAVKKDNSVHWERPQKPKAPVGHFYEPQAPSAEEMTSYVLLAYLTAQPAPTSDELTSATNIV KWITKQNAQGGFSSTQDTVVALHALSKYGAATFTRTGKAAQVTIQSSGTFSSKFQVDNNRLLLQQVSLPELPGEYSM KVTGEGCVYLLQTSCLKYNILPEKEEFPFALGVQTLPTQTCDEPKAHTSFQISLSVSYTGSRASNMAIVDKMVSGFIPLKPTVK MLERSNHVSRTEVSSNHVLIYLDKVSNTLSLFFTVLQDVPVRDLKPAIVKVDYDYETDEFAIAEYNAPCSKDLGNA

靶点信息

研究背景	Alpha-2-macroglobulin is a protease inhibitor and cytokine transporter. It inhibits many proteases, including trypsin, thrombin and collagenase. A2M is implicated in Alzheimer disease (AD) due to its ability to mediate the clearance and degradation of A-beta, the major component of beta-amyloid deposits.
基因ID	2

基因名	A2M
Swiss	P01023
别名	A2M;A2MD;CPAMD5;FWP007;S863-7

产品验证



Western blot analysis of A2M expressed in Mouse uterus, Mouse brain using A2M Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn