

— ABLYBIO, Help Your Research



ZHX2 Rabbit mAb

货号: **AYM29386**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB ICC FC
推荐浓度	WB: 1:500 - 1:2000 ICC: 1:50 - 1:200 FC: 1:20 - 1:50
理论分子量	92kDa
实测分子量	110kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-251MG,293T,Raji,LO2,Mouse brain
细胞定位	cytosol,nucleoplasm,nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human ZHX2.
------	---

靶点信息

研究背景	The members of the zinc fingers and homeoboxes gene family are nuclear homodimeric transcriptional repressors that interact with the A subunit of nuclear factor- κ B (NF- κ B) and contain two C2H2-type zinc fingers and five homeobox DNA-binding domains. This gene encodes member 2 of this gene family. In addition to forming homodimers, this protein heterodimerizes with member 1 of the zinc fingers and homeoboxes family.
基因ID	22882
基因名	ZHX2
Swiss	Q9Y6X8 (https://www.uniprot.org/uniprotkb/Q9Y6X8/entry)
别名	ZHX2,ZHX2 Rabbit mAb,Alpha-fetoprotein regulator 1,Regulator of AFP,Zinc finger and homeodomain protein 2,AFR1,KIAA0854,RAF

产品验证



Western blot analysis of ZHX2 expressed in Mouse brain,U-251MG,293T using ZHX2 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)