

SMARCC1/BAF155 Rabbit mAb

货号: **AYM29389**

产品信息

反应	Human,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IF/ICC IP FC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200 IP: 1:20 - 1:50 FC: 1:20 - 1:50
理论分子量	122kDa
实测分子量	155kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T
细胞定位	Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human SMARCC1/BAF155.
序列	LVVQLLQFQEDAFGKHVTNPAFTKLPACKFMDFKAGGALCHILGAAYKYKNEQGWRFRDLQNP SRMDRNVEMFMNIEKT LVQNNCLTRPNIYLIPDIDLKLANLKLKDIKRHQGTFTDEKSKASHHIYPYSSSQDDEEWLRPVMRKEKQVLVHWGFYPSY DTWVHSNDVDAEIEDPPIPEKPKWKVHVKWLDTDFNEWMNEEDYEVDENRKPVSFRQRISTKNEEPVRSPEPDRKASA

靶点信息

研究背景	The protein encoded by this gene is a member of the SWI/SNF family of proteins, whose members display helicase and ATPase activities and which are thought to regulate transcription of certain genes by altering the chromatin structure around those genes. The encoded protein is part of the large ATP-dependent chromatin remodeling complex SNF/SWI and contains a predicted leucine zipper motif typical of many transcription factors.
------	--

基因ID	6599
基因名	SMARCC1
Swiss	Q92922
别名	SMARCC1;BAF155;CRACC1;Rsc8;SRG3;SWI3

产品验证



Western blot analysis of SMARCC1/BAF155 expressed in 293T using SMARCC1/BAF155 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn