

KGF Rabbit mAb

货号: **AYM29390**

产品信息

| | |
|-------|---|
| 反应 | Human,Mouse,Rat |
| 宿主 | Rabbit |
| 克隆性 | Monoclonal |
| 预测反应 | |
| 应用 | WB |
| 推荐浓度 | WB: 1:500 - 1:2000 |
| 理论分子量 | 23kDa |
| 实测分子量 | 28kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | HeLa,A-549,Raji,Mouse lung,Mouse spleen,Mouse uterus,Rat lung,Rat spleen,Rat uterus |
| 细胞定位 | cytoplasm,extracellular region,extracellular space,Golgi apparatus |
| 纯化 | Affinity purification |

抗原信息


| | |
|------|--|
| 抗原信息 | Recombinant fusion protein corresponding to Human KGF. |
|------|--|

靶点信息

| | |
|------|--|
| 研究背景 | The protein encoded by this gene is a member of the fibroblast growth factor (FGF) family. FGF family members possess broad mitogenic and cell survival activities, and are involved in a variety of biological processes, including embryonic development, cell growth, morphogenesis, tissue repair, tumor growth and invasion. This protein is a potent epithelial cell-specific growth factor, whose mitogenic activity is predominantly exhibited in keratinocytes but not in fibroblasts and endothelial cells. Studies of mouse and rat homologs of this gene implicated roles in morphogenesis of epithelium, reepithelialization of wounds, hair development and early lung organogenesis. [provided by RefSeq, Jul 2008] |
| 基因ID | 2252 |
| 基因名 | FGF7 |

| | |
|-------|--------|
| Swiss | P21781 |
| 别名 | KGF |

产品验证



Western blot analysis of KGF expressed in Mouse lung,Rat lung,HeLa using KGF Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn