

KGF Rabbit mAb

货号: **AYM29390**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	23kDa
实测分子量	28kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,A-549,Raji,Mouse lung,Mouse spleen,Mouse uterus,Rat lung,Rat spleen,Rat uterus
细胞定位	cytoplasm,extracellular region,extracellular space,Golgi apparatus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human KGF.
序列	MHKWILTWILPTLLYRSCFHIICLVGTISLACNDMTPEQMATNVNCSSPERHTRSVDYMEGGDIRVRRLLFCRTQWYLRIKRGKVKGTQEMKNNYNIMEI

靶点信息

研究背景	The protein encoded by this gene is a member of the fibroblast growth factor (FGF) family. FGF family members possess broad mitogenic and cell survival activities, and are involved in a variety of biological processes, including embryonic development, cell growth, morphogenesis, tissue repair, tumor growth and invasion. This protein is a potent epithelial cell-specific growth factor, whose mitogenic activity is predominantly exhibited in keratinocytes but not in fibroblasts and endothelial cells. Studies of mouse and rat homologs of this gene implicated roles in morphogenesis of epithelium, reepithelialization of wounds, hair development and early lung organogenesis. [provided by RefSeq, Jul 2008]
------	--

基因ID	2252
基因名	FGF7
Swiss	P21781
别名	HBGF-7; KGF

产品验证



Western blot analysis of KGF expressed in Mouse lung,Rat lung,HeLa using KGF Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn